

## Immunoglobulins in the Urine of Leprosy Patients

### TO THE EDITOR:

Renal lesions have been noted often in leprosy patients. Most frequently amyloidosis and nephritis of all types, such as chronic pyelonephritis, interstitial nephritis, and chronic diffuse glomerulonephritis, have been reported by various workers (2, 3, 4, 5, 6, 7, 9, 10, 12) in varying frequencies. Although specific leprosy granuloma formation has not been observed in the kidney, it is quite possible that leprosy patients, due to their frequent involvement of renal parenchyma, excrete different components of serum in their urine which directly indicate the extent of renal damage (8). However, the excretion of the type of proteins will depend upon the extent of kidney damage in the leprosy patient. Chakraborty and Saha (1) have noted the presence of IgG and  $\lambda$  chains in the urine of lepromatous leprosy (LL) patients. These authors did not find any serum proteinuria in the tuberculoid type (TT) of leprosy patients.

We tried to demonstrate IgG, IgA, IgM, and C3 in the urine of leprosy patients across the spectrum of the disease. Urine from 26 healthy controls was also taken for comparison for this study. The early morning samples of urine were collected and concentrated ten times before the samples were tested by double diffusion against the specific antisera (anti-immunoglobulins and

anti-C3 were obtained commercially from Kallestad Laboratories, Chaska, Minnesota 55318, U.S.A.). The results are summarized in The Table. Unlike Chakraborty and Saha's (1) findings, 12 healthy control subjects out of 26 showed positivity for  $\kappa$  and 1 out of 26 was also positive for  $\lambda$  chains. However, these samples were negative for the presence of immunoglobulins. Both light chains,  $\kappa$  and  $\lambda$ , are found to be present in varying proportions in all types of leprosy, with 100%  $\kappa$  chain positivity when the patients were in reaction. IgG was present in the urine of all types of leprosy. LL cases with erythema nodosum leprosum (ENL) showed maximum positivity for IgG (75%). IgA was also noted in the urine of all types of leprosy, although in a smaller number of cases. None of the borderline tuberculoid (BT) cases in reaction was positive for IgA. C3 was determined in the urine of nine cases in reaction of BT and ENL types of leprosy. None of these patients showed positivity for C3, although Shwe (11) demonstrated the presence of C3 in renal biopsies of such patients.

—U. Sengupta, Ph.D.

—G. Ramu, M.D.

—Sudhir Sinha, M.Sc.

—V. D. Ramanathan, M.B., B.S., Ph.D.

—K. V. Desikan, M.D.

Central JALMA Institute for Leprosy

P.O. Box 31

Taj Ganj

Agra 282001, India

THE TABLE. *Data showing the presence of light chains, immunoglobulins, and C3 in urine of leprosy patients.*

Source of urine	No.	Light chains positive		Positivity for			
		$\kappa$	$\lambda$	IgG	IgA	IgM	C3
Healthy controls	26	12 <sup>a</sup>	1	0	0	0	0
TT/BT <sup>b</sup>	9	7	3	4	1	0	ND <sup>c</sup>
LL <sup>d</sup>	20	15	16	9	3	0	ND
BT(R) <sup>e</sup>	9	9	5	5	0	0	0
ENL <sup>f</sup>	16	16	13	12	1	0	0

<sup>a</sup> Number of positive cases.

<sup>b</sup> TT = tuberculoid, BT = borderline tuberculoid.

<sup>c</sup> ND = not done.

<sup>d</sup> LL = lepromatous.

<sup>e</sup> R = reaction.

<sup>f</sup> ENL = erythema nodosum leprosum.

### REFERENCES

1. CHAKRABORTY, A. K. and SAHA, K. Immunologic aspects of leprosy with reference to extravascular immunoglobulins. Their excretion profile in urine. *Int. J. Lepr.* **44** (1976) 456-461.
2. DESIKAN, K. V. and JOB, C. K. A review of post-mortem findings in 37 cases of leprosy. *Int. J. Lepr.* **36** (1968) 32-44.
3. GREBSTALD, H. and SWEN, L. Genitourinary lesions in leprosy with reference to the problem of atrophy of testis. *JAMA* **1149** (1952) 1287-1291.
4. GUPTA, J. C., DIWAKAR, R., SINGH, S., GUPTA, D. K. and PANDA, P. K. A histopathologic study of renal biopsies in fifty cases of leprosy. *Int. J. Lepr.* **45** (1977) 167-170.
5. JOHNY, K. V. and KARAT, A. B. A. Renal biopsy

- studies in cases of leprosy. *J. Assoc. Physicians India* **19** (1971) 117-118.
6. KRISHNAMURTHI, S. and JOB, C. K. Secondary amyloidosis in leprosy. *Int. J. Lepr.* **34** (1966) 155-158.
  7. LEDINGHAM, J. M. *The Medical Annual*. Bristol: John Wright and Sons Ltd., 1972, p. 384.
  8. PETERSON, P. A. and GERGGARD, I. Urinary immunoglobulin components in normal tubular and glomerular proteinuria; qualities and characteristics of free light chains, IgG, IgA and Fc fragments. *Eur. J. Clin. Invest.* **1** (1971) 255-273.
  9. POWELL, C. S. and SWAN, L. L. Leprosy: Pathologic changes observed in 50 consecutive necropsies. *Am. J. Pathol.* **31** (1955) 1131-1147.
  10. SATYANARAYANA, B. V., RAJU, P. S., KUMARI, K. R. and REDDY, C. R. R. M. Amyloidosis in leprosy. *Int. J. Lepr.* **40** (1972) 278-280.
  11. SHWE, T. Immune complexes in glomeruli of patients with leprosy. *Lepr. Rev.* **42** (1972) 282-289.
  12. SHUTTLEWORTH, J. S. and ROSS, S. H. Secondary amyloidosis in leprosy. *Ann. Intern. Med.* **45** (1956) 23-28.