

Effect of Preformed Immune Complexes on the Course of *Mycobacterium leprae* Infection in Normal Mice

TO THE EDITOR:

Normal mice infected with *Mycobacterium leprae* in the foot pads show a limited multiplication. Dissemination of infection is seen in immunosuppressed animals, i.e., the thymectomized and irradiated (⁹) and nude mouse (⁷) and the nude rat (⁴). The severe immunosuppression seen in these animals does not simulate human leprosy before or during the course of the disease. Out of the many known and unknown factors operating in the causation of the depressed cell-mediated immunity in lepromatous leprosy, one is the presence of immune complexes (IC). The IC, in turn, are capable of producing specific immunodepression by blocking the recognition of *M. leprae* antigen by lymphocytes. In addition, IC might have multiple other effects on the whole array of immuno-competent cells (²) and, subsequently, dissemination of infection may involve other systems.

In the present study, 4-week-old, closely bred, Swiss albino mice (Lacca strain) were divided into two groups: Group I, composed of 55 normal control mice, and Group II, composed of 55 mice which were inoculated in the right foot pad with *M. leprae* (1×10^4 AFB/foot pad) obtained from biopsy specimens of untreated lepromatous patients. Both the control (NC) and infected (NI) mice were each divided into two batches, i.e., 0dIC (N = 30) and 3mIC (N = 25), depending upon the period at which IC were administered: at zero day (0d) or at 3 months (3m) post-inoculation with *M. leprae*. A

large number of animals were used as a precautionary measure because of their high mortality rate. Five mice each were sacrificed periodically at 3-, 6-, and 9-month intervals from the 0dIC batches and at 6- and 9-month intervals from the 3mIC batches, since the data for the remaining period were obtained from a study conducted and published earlier (¹³) wherein the procedures for foot pad inoculation, harvesting, and counting of bacilli were also mentioned.

Freeze-dried *M. leprae* (obtained as a gift from IMMLEP) were sonicated to prepare antigen (⁶), and healthy rabbits were immunized to raise the anti-*M. leprae* serum (AMLS). The production of antibodies against *M. leprae* antigens was confirmed by immunodiffusion. The zone of equivalence (ZE) was determined as described by Roitt (¹⁰). The tube containing 240 μ g of *M. leprae* antigen gave maximum precipitate and was taken as the ZE. Twice the ZE gave complexes in $2 \times$ antigen excess. AMLS was adsorbed with homogenized human skin tissue and also with *M. bovis* (BCG), *M. tuberculosis* (H37Rv), and *M. avium*. IC were prepared *in vitro* by incubation of the *M. leprae* antigen and the AMLS. IC in $2 \times$ antigen excess (1 mg protein) were administered in 0.3 ml of saline intravenously (i.v.) every week for 1 month starting at day 0 and at 3 months after *M. leprae* inoculation into foot pads.

The kidney, liver, sciatic nerve, skin, earlobes, snout, and tail tip of all sacrificed animals were fixed in 10% Formol saline.




FIG. 1. Kidney section of mouse given human *M. leprae* inoculation and preformed immune complexes. Numerous AFB are seen in the renal parenchyma (Fite-Faraco $\times 330$).




FIG. 2. Liver section of mouse infected with human *M. leprae* and given preformed immune complexes. A pair of AFB is seen in the central vein (Fite-Faraco $\times 330$).

Skin tissue was taken from the right hind leg close to the inoculated site.

In the NI group, data for which were published earlier (¹³), acid-fast bacilli (AFB) recovered from pooled homogenate of five foot pads could not be counted since only a few bacilli were seen in the smears and none were seen in the pin-head stamps. However, there was a steep rise in the AFB counts at 6 months (3.40×10^5 /foot pad) which remained stable up to the 9-month harvests (3.04×10^5 AFB/foot pad). Infection was established early in the NI-0dIC batch, i.e., at 3 months we were able to see bacilli without much difficulty (mean 8.40×10^4 AFB/foot pad). An increased number of AFB could be harvested at the 6-month sacrifice (mean 4.65×10^6 AFB/foot pad) which maintained a plateau up to 9 months (mean 1.79×10^6 AFB/foot pad). Except for the early detection of infection at 3 months, the growth pattern was similar to the NI group. Administration of IC at 3 months did not bring about any appreciable increase in ba-

cillary counts when compared to the NI/0dIC batch at 9 months; counts at the 6- and 9-month harvests were 4.48×10^5 and 5.99×10^5 AFB/foot pad, respectively. No growth was seen in nutrient agar and Löwenstein-Jensen media inoculated with foot pad homogenates.

In the NI group a few AFB were found in 40% of the kidneys at 9 months. Sixty percent of sciatic nerves had a few AFB at 3 and 6 months. At no time were AFB detected in the skin and liver tissues. However, 40% of the animals showed solitary AFB in the earlobes and in only one animal at the nose tip at 9 months.

However, we were surprised to see numerous AFB in parenchyma (Fig. 1) and attached to walls of tubules of 20%, 40%, and 60% of the kidneys at 3-, 6-, and 9-month sacrifices, respectively. The AFB were found in both the cytoplasm of histiocytes and extracellularly. Histopathologically, interstitial nephritis, including periarteritis and periglomerulitis, was also seen

in association with AFB dissemination. A few AFB were detected in 40% and 60% of the sciatic nerves and the skin at 6 and 9 months, respectively. The skin of 20% of the animals at 3 months and the liver (Fig. 2) at each period of study were positive. Focal hepatitis with lymphomononuclear cells and histiocytes were seen in 40% and 20% of the animals at 6 and 9 months, respectively. A few AFB in the tail tip of one animal at both 3 and 6 months and two animals at 9 months were seen. In the NI/3mIC batch, a fairly large number of scattered AFB could be detected in the kidneys (20%) at both harvests. A few bacilli were also seen in 20% of the sciatic nerves and skin. A good number of AFB in the liver and a few AFB in the earlobes, nose tip, and tail tip were seen in one animal each at both 6 and 9 months. In the NC animals, no change was observed in any of the above tissues.

Very little work has been done on the role of IC in the pathogenesis of leprosy in spite of their circumstantial association with erythema nodosum leprosum (ENL) ^(5, 14). Hypergammaglobulinemia ⁽³⁾ results in the combination of specific antibodies with *M. leprae* or host antigens and consequently the production of IC. In the NI group, the infection was established at 6 months and followed a growth curve typical for *M. leprae* infection ⁽¹³⁾. The yields were not as high as obtained by Rees and Weddell ⁽⁹⁾. Variations in the maximum number of AFB obtained and time variables for the infection have been described ⁽¹¹⁾.

AFB could be detected in the various tissues of mice given IC and *M. leprae* inoculation as early as 3 months; whereas in NI mice not administered IC, AFB were seen around 1 year as was reported earlier ⁽¹²⁾. IC seemed to be more effective in the production of immunosuppression when given simultaneously with *M. leprae* inoculation, as seen in the NI/0dIC batch.

From this study it may be envisaged that IC have an immunosuppressive role. Antigenically related mycobacteria may sensitize an individual to *M. leprae* antigens and/or modulate the immune response to a subsequent leprosy infection. If they are related to *M. leprae*, contact even with saprophytic mycobacteria in the environment

might be important ⁽⁸⁾. Clarification of antigenic crossreactivity of *M. leprae* to the other mycobacteria will help in exploring the immuno-epidemiology of leprosy ⁽¹⁾.

—Surrinder Kaur, M.D., M.A.M.S.

—Chetana Vaishnavi, M.Sc., Ph.D.

—Bhushan Kumar, M.D., M.N.A.M.S.

Department of Dermatology

—Nirmal Kumar Ganguly,
M.D., M.N.A.S.

Department of Experimental
Medicine

Postgraduate Institute of
Education and Research
Sector 12

Chandigarh 160012, India

REFERENCES

1. BJUNE, G., CLOSS, O. and BARNETSON, R. St.C. Early events in the host parasite relationship and immune response in clinical leprosy; its possible importance for leprosy control. *Clin. Exp. Immunol.* **54** (1983) 289–297.
2. BULLOCK, W. E. Studies on immune mechanisms in leprosy; depression of delayed allergic response to skin test antigen. *N. Engl. J. Med.* **278** (1968) 298–304.
3. BULLOCK, W. E., HO, M. F. and CHEN, M. J. Studies on immune mechanisms in leprosy; quantitative relationships of IgG, IgA and IgM immunoglobulins. *J. Lab. Clin. Med.* **75** (1970) 863–870.
4. DAWSON, P. J., COLSTON, M. J. and FIELDSTEEL, A. H. Infection of the congenitally athymic rat with *Mycobacterium leprae*. *Int. J. Lepr.* **51** (1983) 336–346.
5. GELBER, R. H., EPSTEIN, V., FASAL, P. and DRUTZ, D. J. Erythema nodosum leprosum, an immune complex disease. *Int. J. Lepr.* **41** (1973) 553–554.
6. HARBOE, M., CLOSS, O., BJORVATN, B., KRONVALL, G. and AXELSEN, N. H. Antibody response in rabbits to immunization with *Mycobacterium leprae*. *Infect. Immun.* **18** (1977) 792–805.
7. LANCASTER, R. D., HILSON, G. R. F., McDougall, A. C. and COLSTON, M. J. *Mycobacterium leprae* infection in nude mice; bacteriological and histological responses to primary infection and large inocula. *Infect. Immun.* **39** (1983) 865–872.
8. PAUL, R. C., STANFORD, J. L. and CARSWELL, J. W. Multiple skin testing in leprosy. *J. Hyg. (Camb.)* **75** (1975) 57–68.
9. REES, R. J. W. and WEDDELL, A. G. M. Experimental models for studying leprosy. *Ann. N.Y. Acad. Sci. U.S.A.* **154** (1968) 214–236.
10. ROITT, I. M. The classical precipitin reaction. In: *Essential Immunology*. 4th ed. London: Blackwell Scientific, 1980, pp. 5–6.

11. SHEPARD, C. C. and McRAE, P. H. Hereditary characteristic that varies among isolates of *Mycobacterium leprae*. *Infect. Immun.* **3** (1971) 121–126.
12. VAISHNAVI, C., KAUR, S., KUMAR, B., GANGULY, N. K. and CHAKRAVARTI, R. N. Trophic changes and extensive dissemination in normal mice infected with human *Mycobacterium leprae*. *Indian J. Lepr.* **56** (1984) 742–747.
13. VAISHNAVI, C., KUMAR, B., GANGULY, N. K. and KAUR, S. Phagocytic and bactericidal activities of macrophages from *Mycobacterium leprae* infected normal and immunosuppressed mice. *Int. J. Lepr.* **53** (1985) 233–237.
14. WEMAMBU, S. N. C., TURK, J. L., WATERS, M. F. R. and REES, R. J. W. Erythema nodosum leprosum: a clinical manifestation of the Arthus phenomenon. *Lancet* **2** (1969) 933–935.