

Roles of Tumor Necrosis Factor- α and Transforming Growth Factor- β in Regulating Intercellular Adhesion Molecule-1 Expression on Murine Peritoneal Macrophages Infected with *M. leprae*¹

Toshiaki Shimizu, Win Win Maw, and Haruaki Tomioka²

Adhesion molecules expressed on immunocompetent cells play important roles in cellular interactions for the development of immunological responses (²⁴). Intercellular adhesion molecule-1 (ICAM-1) plays a role in lymphocyte trafficking by allowing ICAM-1-bearing endothelial cells to interact with leukocyte function-associated antigen-1 (LFA-1) on lymphocytes, and thereby facilitate the migration of lymphocytes to sites of inflammation along endothelial cells (²⁴). Moreover, the interaction of LFA-1 with ICAM-1 is critical for effective cellular interactions of T cells with antigen-presenting cells, including macrophages (M Φ s), and the consequent activation of resting T cells (^{12, 15, 24}). Indeed, ICAM-1 augments T-cell proliferation and is pivotal in recall antigen responses of T cells to purified protein derivative of *Mycobacterium tuberculosis* (³⁰). Moreover, a recent study of Lopez Ramirez, *et al.* (¹⁴) showed that the ICAM-1 expression on THP-1 human monocyte-derived cell line was potently increased by stimulation with *M. tuberculosis*. In addition, the *in vivo* study by Sullivan, *et al.* (²⁵) concerning *M. leprae* infection revealed that granulomas from both tuberculoid leprosy and lepromatous leprosy displayed ICAM-1 expression. This implies that ICAM-1 plays important roles in enhancing the immune and inflammatory

responses to *M. leprae*. Therefore, it is of interest to know the profiles of ICAM-1 expression by M Φ s after stimulation with *M. leprae*. In the present study, we found that ICAM-1 expression on murine peritoneal M Φ s was increased during days 1 to 3 after stimulation with *M. leprae* because of autocrine upregulation by tumor necrosis factor-alpha (TNF- α). The ICAM-1 expression thereafter gradually decreased, returning to the normal level by day 7 due to the downregulating effect of transforming growth factor-beta (TGF- β) derived from *M. leprae*-stimulated M Φ s themselves.

MATERIALS AND METHODS

Organisms. *M. leprae* were purified from the liver of an infected armadillo (#825) obtained from G. P. Walsh, Armed Forces Institute of Pathology, Washington, DC, U.S.A., by Percoll gradient centrifugation according to the method of Draper (IMMLEP protocol 1/79) with slight modifications (²¹). This *M. leprae* strain 825 contained 30% of living cells on the basis of fluorescein diacetate and ethidium bromide (FDA/EB) staining and 10⁹ of the organisms contained 7.4 pmoles of ATP, indicating good viability of the bacterial preparation. In some experiments, *M. leprae* organisms were collected and purified from infected foot pads of BALB/c nude mice, as previously reported (²⁶). When 10⁸ of these nude mouse *M. leprae* (strain Thai-53) were inoculated in the 7H12B medium of the BACTEC 460 Tuberculosis System and cultured according to Franzblau (¹¹), the growth index (GI) value increased from 0 to 122 and 267 at day 4 and day 11, respectively, indicating good viability of this bacterial preparation. When BALB/c nude

¹ Received for publication on 25 November 1997. Accepted for publication in revised form on 10 December 1998.

² T. Shimizu, M.S.; W. W. Maw, M.D., Ph.D.; H. Tomioka, Ph.D., Department of Microbiology and Immunology, Shimane Medical University, Izumo, Shimane 693, Japan.

Reprint requests to Dr. H. Tomioka at the above address or FAX 81-853-20-2145.

mice were infected with 10^6 of *M. leprae* Thai-53 into the hind foot pad, the organisms steadily grew at the sites of infection, reaching bacterial loads larger than 10^9 bacilli per foot pad within 1 year after infection (data not shown). This also indicates good viability of the bacterial preparation used.

Mice. Seven- to ten-week-old, male BALB/c mice purchased from Japan Clea Co., Osaka, Japan, were used.

Special agents for M Φ culture. Recombinant mouse TNF- α , ultrapure natural human TGF- β 1 and mouse anti-human TGF- β monoclonal antibody (mAb) (also specific to mouse TGF- β) were purchased from Genzyme Co., Cambridge, Massachusetts, U.S.A. Rat anti-mouse TNF- α mAb was obtained from Pharmingen Co., San Diego, California, U.S.A. One unit of TNF- α is defined as the amount required to mediate half-maximal cytotoxicity of L929 cells. All of these agents contained no preservative, such as sodium azide, and were found to be essentially free from LPS contamination by the Limulus J Single Test (Wako Pure Chemical Industry Co., Osaka, Japan).

Medium. Medium RPMI 1640 (Nissui Pharmaceutical Co., Tokyo, Japan) supplemented with 25 mM HEPES, 2 mM glutamine, and 5% heat-inactivated fetal bovine serum (FBS) (M. A. Bioproducts, Walkersville, Maryland, U.S.A.) was used for cell culture unless otherwise specified. The medium and FBS used were found to be free from LPS contamination by the Limulus J Single Test.

Peritoneal M Φ s. Peritoneal exudate cells (PECs) were collected with Hanks' balanced salt solution (HBSS) containing 2% FBS from mice given an intraperitoneal injection of 1 ml of peptone-starch 4 days before harvest. After washing with 2% FBS-HBSS by centrifugation at $200 \times g \times 5$ min, the PECs were suspended in a small volume of the culture medium. Ten ml each of PEC suspension in 5% FBS-RPMI 1640 at the cell density of 5×10^6 /ml was poured onto a 90-mm cell culture plate (Corning Glass Works Co., Corning, New York, U.S.A.) which was overlaid with 14-mm plastic sheets (about 20 sheets) (Wako Pure Chemical Industry Co., Osaka, Japan). After a 2-hr incubation at 37°C in a CO₂ incu-

bator (5% CO₂-95% humidified air), the resultant sheets were removed, thoroughly rinsed with 2% FBS-HBSS, and immersed into 1 ml of culture medium in a 16-mm cell culture well (24-wells; Corning). This method gave more than 90% pure M Φ cultures, with potent pinocytic ability for neutral red and phagocytic activity for latex beads.

Microscopic counting for ICAM-1-positive M Φ s. M Φ monolayer cultures on the plastic sheets were cultured in 1.0 ml of 5% FBS-RPMI 1640 in 16-mm culture wells (Corning) in the presence or absence of 1×10^7 /ml of *M. leprae* or latex beads (3.0- μ m diameter; Sigma Chemical Co., St. Louis, Missouri, U.S.A.) at 37°C in a CO₂ incubator for up to 14 days. At intervals, the M Φ monolayer cultures were removed, washed with phosphate buffered saline (PBS) containing 0.2% bovine serum albumin (BSA), fixed with 1% paraformaldehyde, incubated in 1% BSA-PBS at room temperature for 30 min, and then reacted with rat anti-mouse ICAM-1 mAb (Seikagaku Corp., Tokyo, Japan) for 1 hr. After rinsing with 0.1% BSA-PBS, the M Φ s were reacted with alkaline phosphatase-conjugated anti-rat Ig mAb (Seikagaku) for 1 hr and then washed with 0.1% BSA-PBS. Color development was achieved by using NBT-BCIP substrate (nitroblue tetrazolium/5'-bromo-4-chloro-3-indolylphosphate) and the proportion of blue-stained M Φ s (ICAM-1-positive cells) was enumerated by microscopic counting (¹⁶).

Cytokine measurement. The 1-, 3- or 7-day culture fluids of M Φ s with or without *M. leprae* infection were measured for TNF- α and TGF- β concentrations as previously described. Briefly, Immulon 4 plates (Dynatech Laboratories, Chantilly, Virginia, U.S.A.) were coated with a capture Ab for each cytokine using rat anti-mouse TNF- α mAb (Pharmingen) and mouse anti-human TGF- β mAb (also specific to mouse TGF- β) (Genzyme). Biotinylated rat anti-mouse TNF- α mAb (Pharmingen) and chicken anti-human TGF- β mAb (R & D System, Inc., Minneapolis, Minnesota, U.S.A.) were used as the detecting Abs. After binding of alkaline phosphatase-conjugated streptavidin (Life Technologies Co., Gaithersburg, Maryland, U.S.A.) in the case of TNF- α assay and alkaline phosphatase-




FIG. 1. Profiles of ICAM-1 expression by *M. leprae* 825-infected MΦs during cultivation in medium with or without addition of exogenous TNF- α . \square = Uninfected MΦs were cultured in the absence of TNF- α ; ▨ = *M. leprae*-infected MΦs were cultured in the absence of TNF- α ; \blacksquare = *M. leprae*-infected MΦs were cultured in the presence of TNF- α added at 500 units/ml on day 0. Each bar indicates mean \pm S.E.M. (N = 2).

conjugated rabbit anti-chicken/turkey IgG Ab (Zymed Laboratories Inc., San Francisco, California, U.S.A.) in the case of TGF- β assay, color development was achieved using *p*-nitrophenyl phosphate tablets (Sigma) as the substrate. Capture Ab and detecting Ab for each cytokine had different epitope specificities from each other. The present ELISA method was capable of measuring the active form TGF- β in MΦ culture fluids in a specific fashion.

RESULTS

ICAM-1 expression on *M. leprae*-infected MΦs during cultivation. As shown in Figure 1, when MΦs were infected with *M. leprae* at day 0 (hatched bars), the ICAM-1 expression of cultured MΦs, in terms of increase in the ratio of ICAM-1-positive MΦs measured by the microscopic method, rapidly increased reaching the peak in the early phase (during day 1 to day 3) of cultivation and thereafter decreased, returning to the normal level by day 7. A lower increase in the ICAM-1 expression was also noted for uninfected (unstimulated) MΦs in the same period of time (open bars). In this experiment, residual numbers of attached MΦs on a plastic culture sheet during cultivation after *M. leprae*




FIG. 2. Effects of exogenous TNF- α on ICAM-1 expression by uninfected MΦs during cultivation. MΦs were cultivated in the presence (\blacksquare) or absence (\square) of TNF- α which was added at 500 units/ml on days 0, 2, and 6 during cultivation. Each bar indicates mean \pm S.E.M. (N = 3). * Significantly larger than value of MΦs cultured in absence of TNF- α ($p < 0.05$; Student's *t* test).

rae-infection (hatched bars) were changed as follows when the day 0 value was fixed as 100% (N = 5): day 1, $97.6 \pm 5.5\%$; day 3, $91.0 \pm 7.1\%$; day 7, $78.9 \pm 6.9\%$. This indicates that the reduction of ICAM-1 expression in the middle phase (day 7) of MΦ cultivation after *M. leprae*-infection was not due to *M. leprae*-mediated cytotoxicity. Figure 1 also shows that the addition of TNF- α at the concentration of 500 units/ml overcame the middle phase (day 7) downregulation of the ICAM-1 expression (shaded bar), indicating that ICAM-1 downregulation in the middle phase was due in part to a defect of TNF- α . In this context, separate experiments showed that TNF- α , a crucial ICAM-1 upregulating cytokine^(11,20) displayed its ICAM-1 upregulating effect on MΦs as follows.

First, as shown in Figure 2, a marked increase in the ICAM-1 expression was also observed for uninfected MΦs, when treated with exogenous TNF- α (500 units/ml), which was added on days 0, 2, and 6, in the early phase of cultivation (days 1 to 3) but not in the middle phase (day 7). Thus, in the case of uninfected MΦs, exogenous TNF- α could increase only their early phase ICAM-1 expression, differing from the case of *M. leprae*-infected MΦs (Fig. 1). Second, as shown in Figure 3, the addition of




FIG. 3. Effects of anti-TNF- α Ab on ICAM-1 expression by *M. leprae* Thai-53-infected M Φ s. \square = Uninfected M Φ s were cultured in the absence of anti-TNF- α Ab; \square with diagonal lines = *M. leprae*-infected M Φ s were cultured in the absence of anti-TNF- α Ab; \blacksquare = *M. leprae*-infected M Φ s were cultured in the presence of anti-TNF- α Ab which was added at 10 μ g/ml to the culture medium on days 0, 2, and 6 during cultivation. Each bar indicates mean \pm S.E.M. (N = 3). * Significantly smaller than the value of M Φ s cultured in the absence of anti-TNF- α Ab ($p < 0.05$; Student's *t* test).

anti-TNF- α Ab (10 μ g/ml) caused the reduction of the ICAM-1 expression by *M. leprae*-infected M Φ s and this effect was most marked at day 3, confirming that M Φ -derived endogenous TNF- α plays the most important role in the early phase (around day 3) increase in the ICAM-1 expression by *M. leprae*-infected M Φ s.

The above findings indicate central roles of TNF- α in ICAM-1 upregulation of *M.*

leprae-infected M Φ s. However, as indicated in The Table only small amounts of TNF- α were released from *M. leprae*-infected M Φ s into the culture medium during cultivation periods for up to 7 days. It thus appears that the majority of the TNF- α molecules were produced by *M. leprae*-infected M Φ s in a membrane-bound form, molecular weight (MW) of 26 kD, but not in a released form (MW of 17 kD) (¹³). This membrane-bound form of TNF- α seems to play a crucial role in the ICAM-1 upregulation of *M. leprae*-infected M Φ s.

Another important finding in Figure 1 is that TNF- α failed to overcome the late phase (day 14) downregulation of ICAM-1 expression. This phenomenon may be explained as follows. First, ICAM-1 reduction in the late phase seems to be due in part to nutrient deficiency in the M Φ culture medium, as previously demonstrated for *M. avium* complex-infected M Φ s (¹⁶). Second, it is plausible that, in the middle-to-late phase of cultivation, *M. leprae*-infected M Φ s produced certain factors possessing an ICAM-1 downregulatory activity, as described below.

Role of TGF- β in the middle-to-late phase downregulation of ICAM-1 expression by *M. leprae*-infected M Φ s. M Φ s stimulated with mycobacterial organisms produce TGF- β , an immunosuppressive and M Φ -deactivating cytokine (^{4, 31}). We previously found that the *in vitro* TGF- β production by murine peritoneal M Φ s was initiated after day 3 and continued for up to

THE TABLE. Production of TNF- α and TGF- β by *M. leprae*-infected M Φ s during cultivation in the presence or absence of anti-TNF- α or anti-TGF- β Ab.^a

<i>M. leprae</i> infection	Addition of Ab	TNF- α production (pg/ml) ^b			TGF- β production (pg/ml) ^b		
		Day 1	Day 3	Day 7	Day 1	Day 3	Day 7
—	—	20 \pm 2	20 \pm 10	20 \pm 10	50 \pm 10	360 \pm 10	330 \pm 10
+	—	— ^c	10 \pm 8	40 \pm 10	50 \pm 10	490 \pm 10 ^d	420 \pm 30
+	Anti-TNF- α	ND ^e	ND	ND	50 \pm 10	640 \pm 10 ^f	600 \pm 20 ^f
+	Anti-TGF- β	— ^c	— ^c	40 \pm 2	ND	ND	ND

^a M Φ s infected with *M. leprae* Thai-53 were cultured in medium with or without addition of indicated anti-cytokine Abs. Ten μ g each of anti-TNF- α or anti-TGF- β Ab was added to the culture medium (1.0 ml) on days 0, 2, and 6. At intervals, culture fluids were measured for TNF- α and TGF- β concentrations by ELISA.

^b The mean \pm S.E.M. (N = 3).

^c Lower than the detectable limit for TNF- α (10 pg/ml) or TGF- β (10 pg/ml).

^d Significantly larger than the value of uninfected M Φ s ($p < 0.005$; Student's *t* test).

^e ND = Not determined.

^f Significantly larger than the value of *M. leprae*-infected M Φ s cultured in the absence of anti-cytokine Ab (^f $p < 0.005$; ^g $p < 0.05$; Student's *t* test).




FIG. 4. Effects of anti-TGF- β Ab on ICAM-1 expression by *M. leprae* 825-infected M Φ s. \square = Uninfected M Φ s were cultured in the absence of anti-TGF- β Ab; ▨ = *M. leprae*-infected M Φ s were cultured in the absence of anti-TGF- β Ab; \blacksquare = *M. leprae*-infected M Φ s were cultured in the presence of anti-TGF- β Ab which was added at 30 $\mu\text{g/ml}$ to the culture medium on days 0, 3, and 6 during cultivation. Each bar indicates mean \pm S.E.M. (N = 3). * Significantly larger than value of M Φ s cultured in absence of anti-TGF- β Ab ($p < 0.05$; Student's *t* test).

at least day 14 when the M Φ s were infected with *M. avium* complex (¹⁶). As indicated in The Table, significant levels of TGF- β were produced into the culture medium by M Φ s regardless of *M. leprae* infection. However, it was also found that *M. leprae*-infected M Φ s produced significantly larger amounts of TGF- β than did uninfected M Φ s. Notably, anti-TNF- α Ab enhanced the TGF- β production by *M. leprae*-infected M Φ s. In a separate experiment, the addition of 20 $\mu\text{g/ml}$ of anti-TGF- β Ab caused an $82 \pm 2\%$ reduction of TGF- β accumulation into culture fluids of *M. leprae*-infected M Φ s on day 3.

In addition to these findings, it has also been reported that TGF- β displayed down-regulatory effects on the interferon-gamma (IFN- γ)-induced ICAM-1 expression by rat microglia cells, which are cells of M Φ lineage (³³). It is thus likely that TGF- β also plays some active role in ICAM-1 down-regulation observed in the case of *M. leprae*-infected M Φ s. As shown in Figure 4, anti-TGF- β Ab significantly blocked the middle phase (day 7) down-regulation of ICAM-1 expression by M Φ s infected with *M. leprae*. In this case, anti-TGF- β Ab was




FIG. 5. Effects of anti-TGF- β Ab on ICAM-1 expression by M Φ s stimulated with heat (121°C, 15 min)-killed *M. leprae* 825. \square = ICAM-1 expression by normal M Φ s before stimulation with heat-killed *M. leprae*; ▨ = *M. leprae*-stimulated M Φ s were cultured in the absence of anti-TGF- β Ab; \blacksquare = *M. leprae*-stimulated M Φ s were cultured in the presence of anti-TGF- β Ab which was added at 30 $\mu\text{g/ml}$ to culture medium on days 0, 3, and 6 during cultivation. Each bar indicates mean \pm S.E.M. (N = 4). * Significantly larger than value of M Φ s cultured in absence of anti-TGF- β Ab ($p < 0.05$; Student's *t* test).

added at 30 $\mu\text{g/ml}$ on days 0, 3, and 6. In separate experiments, anti-TGF- β Ab at 30 $\mu\text{g/ml}$ was found to neutralize 12 ng/ml of TGF- β , which is much larger than the amounts of TGF- β produced by *M. leprae*-infected M Φ s (The Table).

Figure 5 shows the effects of anti-TGF- β Ab on the ICAM-1 expression by M Φ s stimulated with heat-killed (121°C, 15 min) *M. leprae*. In this case, anti-TGF- β Ab similarly blocked the middle phase (day 7) down-regulation of ICAM-1, thereby indicating that profiles of ICAM-1 expression in M Φ s stimulated due to infection with living *M. leprae* organisms were essentially the same as those in M Φ s stimulated with heat-killed *M. leprae*.

Effects of addition of TNF- α or anti-TGF- β Ab on ICAM-1 expression by latex beads-phagocytosing M Φ s. In order to know whether or not TNF- α -mediated up-regulation and TGF- β -mediated down-regulation of M Φ ICAM-1 expression observed in Figures 1, 4, and 5 was specifically caused by *M. leprae* infection, we examined the effects of the addition of TNF- α and anti-TGF- β Ab on the ICAM-1 expression by M Φ s stimulated by phagocytosis of latex beads. For this purpose, M Φ s were




FIG. 6. Profiles of ICAM-1 expression by latex beads-phagocytizing MΦs during cultivation. □ = Day 0: ICAM-1 expression by normal MΦs before latex phagocytosis. Days 1, 3, 7: ICAM-1 expression by control MΦs without phagocytizing latex particles after cultivation in medium without addition; ▨ = latex beads-phagocytizing MΦs cultured in medium without addition; ■ and ▩ = latex beads-phagocytizing MΦs cultured in presence of either TNF- α (■) or anti-TGF- β Ab (▩). TNF- α and anti-TGF- β Ab added to culture medium at 500 units/ml and 10 μ g/ml, respectively, on days 0, 2, 6 during cultivation. Each bar indicates mean \pm S.E.M. (N = 3). * Significantly larger than the value of control MΦs (p < 0.05; Student's *t* test).

cultured with 3- μ m latex beads which are known to be taken up by MΦs through phagocytosis. As shown in Figure 6, the ICAM-1 expression by latex beads phagocytizing MΦs did not reduce so markedly at day 7 as in the case of *M. leprae*-infected MΦs (Figs. 1, 3, and 4). Moreover, neither TNF- α nor anti-TGF- β Ab significantly affected the ICAM-1 expression by latex beads phagocytizing MΦs.

DISCUSSION

The present study indicated that stimulation of murine peritoneal MΦs with *M. leprae* caused an upregulation of ICAM-1 expression by the MΦs in the early phase (days 1 to 3) of cultivation. However, the ICAM-1 expression thereafter rapidly decreased, and returned to the normal level by day 7. This profile of change in MΦ ICAM-1 expression during the course of cultivation seems to be peculiar to *M. leprae*-infected MΦs, since such marked decrease in the ICAM-1 expression at day 7 was not observed for MΦs stimulated due to phagocytosis of latex beads (Fig. 6). The middle phase (day 7) downregulation of

ICAM-1 expression appeared to be due in part to a defect of MΦ-derived TNF- α having upregulating effects on MΦ ICAM-1 expression (^{14, 20}), since supplementary addition of TNF- α prevented the middle (day 7) phase reduction of ICAM-1 expression (Fig. 1).

This concept is supported by the following findings obtained in separate experiments. First, exogenous TNF- α increased the ICAM-1 expression on uninfected MΦs (Fig. 2). Second, anti-TNF- α Ab abolished the *M. leprae* infection-induced increase in MΦ expression of TNF- α in the early phase (day 3) of cultivation (Fig. 3). However, under the present experimental condition, we detected only low levels of TNF- α released from *M. leprae*-infected MΦs into the culture medium (The Table). This may mean the individual *M. leprae*-infected MΦs produced TNF- α mainly in a membrane-bound form with a MW of 26 kD but not in a releasing form with a MW of 17 kD (^{13, 27}), and that such membrane-bound TNF- α molecules transfer their stimulatory signals to other MΦs through cell-to-cell contact. Alternatively, it is also possible that MΦ-derived endogenous TNF- α could induce the ICAM-1 upregulation of *M. leprae*-infected MΦs even at very low concentrations.

It appears that an immunoregulatory cytokine, TGF- β (³¹) also plays an important role in the ICAM-1 downregulation of the MΦs, since anti-TGF- β Ab overcame the middle phase (day 7) downregulation of the ICAM-1 expression by MΦs infected or stimulated with *M. leprae* (Figs. 4 and 5). In this context, anti-TGF- β Ab failed to increase the TNF- α released by *M. leprae*-infected MΦs (The Table). Therefore, TGF- β -mediated downregulation of MΦ ICAM-1 expression is not attributable to the reduction of MΦ TNF- α -producing ability due to TGF- β action. Moreover, although a significant level of the TGF- β production by *M. leprae*-infected MΦs was observed at day 3, ICAM-1 downregulation was not observed at this time point (The Table versus Figs. 1 and 4). This suggests that TGF- β could not effectively suppress the ICAM-1 expression by MΦs which were actively producing ICAM-1 upregulating factors, including TNF- α . Moreover, it seems that MΦ factors other than TGF- β

also contribute to the middle-to-late phase downregulation of the ICAM-1 expression. For instance, the antiinflammatory cytokine interleukin 10 (IL-10) (^{8, 19, 22}) and, also, prostaglandin E₂ may play some roles in the ICAM-1 downregulation. Indeed, our recent study on *M. avium* complex-infected MΦs revealed that the middle (day 7) to late (day 14) phase ICAM-1 expression by *M. avium* complex-infected MΦs is potently suppressed by these MΦ factors in an autocrine fashion (Tomioka, *et al.*: manuscript submitted). Thus, it is of interest to examine the participation of these and other unknown ICAM-1 downregulating factors in the late-phase reduction of MΦ ICAM-1 expression, with special reference to their collaborating effects.

It is somewhat enigmatic to note that both exogenous TNF-α and anti-TGF-β Abs fully overcame the middle phase (day 7) downregulation of ICAM-1 expression on *M. leprae*-infected MΦs (Figs. 1 and 4). Although the precise reason for this situation is unknown, it may be explained as follows. First, it is thought that by receiving high levels of activating signals of TNF-α MΦs might become resistant to ICAM-1 expression downregulatory effects of TGF-β. If this was the case, the addition of exogenous TNF-α in excess amounts might cause the effective recovery of MΦ ICAM-1 expression in the middle phase (day 7) by protecting *M. leprae*-infected MΦs from TGF-β-mediated ICAM-1 downregulating effects despite the fact that significantly high amounts of TGF-β were produced and released by the MΦs into the culture medium at the same time point. Second, it is also thought that TNF-α-mediated activating signals suppress TGF-β producing ability of *M. leprae*-infected MΦs, since anti-TNF-α Ab was found to decrease MΦ TGF-β production (The Table). Therefore, TNF-α treatment of *M. leprae*-infected MΦs might antagonize TGF-β-mediated ICAM-1 downregulation by reducing the production of MΦ TGF-β, causing a recovery of MΦ ICAM-1 expression.

Notably, the profiles of ICAM-1 expression in MΦs stimulated with living *M. leprae* organisms were essentially the same as those in MΦs stimulated with heat-killed bacilli (Figs. 4 and 5). This indicates that heat-stable components of *M. leprae* play

roles not only in the early-phase upregulation ICAM-1 expression but also in the middle-to-late phase downregulation of ICAM-1 expression. First, lipoarabinomannan (LAM) is one of the major candidates for the ICAM-1-modulating components of *M. leprae*, since mycobacterial LAM is capable of upregulating MΦ ICAM-1 expression (¹⁴). It has been reported that arabinosylated LAM of *M. tuberculosis* induces MΦ production of proinflammatory cytokines, including TNF-α and IL-1 (^{2, 5, 7, 18}), which are the major mediators of ICAM-1 upregulation (^{10, 14, 20}). Nevertheless, in the present study, *M. leprae*-infected MΦs did not produce detectable amounts of TNF-α in the culture medium (The Table). In this context, Adams, *et al.* (¹) reported that LAM from *M. leprae* having mannose capping did not induce TNF-α production by MΦs, as in the case of mannosylated LAM from virulent *M. tuberculosis* Erdman strain (LAM_{Erdman}) (⁶). However, they also found that mannosylated LAM_{Erdman} could trigger MΦ antimicrobial activity and elevated levels of MΦ nitric oxide production, and that this triggering capacity of LAM_{Erdman} was abrogated by an anti-TNF-α Ab, indicating that TNF-α is involved in the induction of MΦ activation by LAM_{Erdman}. Notably, LAM_{Erdman} did so without inducing elevated levels of TNF-α in the supernatant medium. As discussed above, it thus appears that the LAM of *M. leprae* triggered MΦs to produce TNF-α in a membrane-bound form (MW = 26 kD) (¹³), and such membrane-bound TNF-α molecules transferred their stimulatory signals from one MΦ to another MΦ through cell-to-cell contact. Indeed, it has been demonstrated that activated MΦs could kill a TNF-α-sensitive tumor cell line in the absence of any demonstrable TNF-α in the culture supernatants and this killing was attributed to the presence of cell surface-associated TNF-α (⁹). In addition, segments of *M. leprae* cell-wall skeleton, such as mycolyl-arabinogalactan-peptidoglycan complex, protein-peptidoglycan complex, and muramyl dipeptide, may play roles in the upregulation of ICAM-1 expression by *M. leprae*-infected MΦs since these substances are known to trigger elevated production of TNF-α (³).

Second, mycobacterial LAM also induces M Φ production of immunoregulatory cytokines, including TGF- β and IL-10^(7,29) which have ICAM-1 downregulating functions^(23,32), suggesting the roles of LAM in the ICAM-1 downregulation of *M. leprae*-infected M Φ s in the middle-to-late phase of cultivation. In addition, it is also likely that other cellular components, such as heat-stable proteins, play roles in the ICAM-1 modulation since purified protein derivative (PPD) of *M. tuberculosis* is reported to upregulate M Φ production of TNF- α , IL-1, and, moreover, TGF- β ^(10,29).

Moncada, *et al.*⁽¹⁷⁾ reported that the epidermis of leprosy patients with inflammatory skin diseases exhibited an increased ICAM-1 expression. They also found a lack of ICAM-1 expression in the epidermis, especially keratinocytes (cells with M Φ -like functions), of lepromatous patients. It is of interest to know whether or not the defective expression of ICAM-1 in the lesional epidermis of lepromatous leprosy patients is mediated by TGF- β . In this context, we previously found sequential increase of tissue levels of TNF- α , IL-10 and TGF- β in spleens during the course of *M. avium* complex infection in mice⁽²⁸⁾. That is, the increase in the TNF- α level was observed around week 2 after infection, followed by an elevated level of IL-10 during weeks 2 to 4, and the increase of TGF- β level was subsequently observed after week 4 for at least up to week 8. Further studies are currently underway to examine the roles of ICAM-1 downregulating cytokines, including TGF- β and IL-10 as mediators of defective ICAM-1 expression in skin lesions of lepromatous leprosy patients.

SUMMARY

Profiles of intercellular adhesion molecule-1 (ICAM-1) expression on murine peritoneal macrophages (M Φ s) infected with *Mycobacterium leprae* during cultivation were examined with special reference to the regulatory effects of tumor necrosis factor-alpha (TNF- α) and transforming growth factor-beta (TGF- β). When M Φ s were infected with *M. leprae* or stimulated with heat-killed *M. leprae* at day 0, their ICAM-1 expression, measured in terms of the ratio of M Φ s positively stained with anti-ICAM-1 antibody (Ab), rapidly in-

creased, peaking during days 1 to 3 and thereafter fell, returning to the normal level by day 7. The addition of TNF- α or anti-TGF- β Ab inhibited the middle phase (day 7) downregulation of M Φ ICAM-1 expression, although the late-phase (day 14) downregulation of ICAM-1 was not prevented by them. *M. leprae*-infected M Φ s released small amounts of TNF- α and significant amounts of TGF- β into the culture medium. This may indicate that *M. leprae*-infected M Φ s produced the majority of TNF- α in a membrane-bound form. Alternatively, endogenous TNF- α might upregulate M Φ ICAM-1 expression even at very low concentrations. In any case, these findings indicate the central roles of TNF- α and TGF- β in the early phase upregulation and the middle-to-late phase downregulation, respectively, of ICAM-1 expression by *M. leprae*-infected M Φ s.

RESUMEN

Se examinó el efecto regulatorio del factor de necrosis tumoral alfa (TNF α) y del factor de crecimiento transformante beta (TGF β) sobre la expresión de la molécula de adhesión intercelular ICAM-1 en macrófagos peritoneales murinos infectados con *Mycobacterium leprae*.

Los macrófagos infectados con *M. leprae* o estimulados con *M. leprae* muerto por calor en el día 0, mostraron un rápido incremento en la expresión de ICAM-1 que fue máxima entre los días 1-3 y luego decayó, retornando a su nivel normal hacia el día 7. La adición de TNF α o de un anticuerpo anti-TGF- β inhibió la fase media (día 7) de la expresión de ICAM-1 pero no modificó la fase tardía (día 14). Los macrófagos infectados con *M. leprae* liberaron pequeñas cantidades de TNF α y cantidades significativas de TGF- β en el medio de cultivo. Esto puede indicar que los macrófagos infectados por *M. leprae* produjeron la mayor parte del TNF α en una forma unida a membranas.

Alternativamente, el TNF α endógeno podría regular la expresión de ICAM-1 aun a muy bajas concentraciones. En cualquier caso, estos hallazgos indican que el TNF- α y el TGF- β participan en la regulación de la expresión de ICAM en las fases temprana y tardía, respectivamente, de la infección por *M. leprae*.

RÉSUMÉ

On a examiné le mode d'expression des molécules d'adhésion intercellulaire de type 1 (ICAM-1) chez des macrophages (M Φ s) infectés *in vitro* par *Mycobacterium leprae*, en tenant compte en particulier de la régulation d'expression exercée par le tumor necrosis factor-alpha (TNF- α) et le transforming growth factor-beta (TGF- β).

Lorsque des MΦs étaient infectés par *M. leprae* ou bien stimulés par des *M. leprae* tuées par la chaleur, l'expression de ICAM-1, mesurée en terme de proportion de MΦs marqués par des anticorps (Ac) anti-ICAM-1, augmentait rapidement, présentait un pic entre le premier et le troisième jour après l'infection et diminuait ensuite, pour rejoindre le niveau basal d'expression vers le septième jour. L'addition de TNF- α ou d'anticorps anti-TGF- β inhibait la phase intermédiaire (septième jour) de retour au niveau basal d'expression d'ICAM-1 par les MΦs. Cependant, le retour tardif (quatorzième jour) au niveau de base d'expression de ICAM-1 n'a pu être prévenu par ces molécules. Les MΦs infectés par *M. leprae* ont relargués dans le milieu de culture une petite quantité de TNF- α et une quantité plus importante de TGF- β .

Cela pourrait suggérer que soit les MΦs infectés par *M. leprae* produiraient la majorité de leur TNF- α dans une forme associée aux membranes, soit que le TNF- α endogène pourrait stimuler l'expression de ICAM-1 même à très petite concentration chez les macrophages. En tout état de cause, ces trouvailles suggèrent un rôle central de TNF- α et de TGF- β , dans la stimulation précoce et dans le retour au niveau basal en phase moyenne et tardive, respectivement, de l'expression de ICAM-1 par les MΦs infectés par *M. leprae*.

Acknowledgment. This study was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, and Culture of Japan, and by the United States-Japan Cooperative Medical Science Program (Tuberculosis and Leprosy Section).

REFERENCES

- ADAMS, L. B., FUKUTOMI, Y. and KRAHENBUHL, J. L. Regulation of murine macrophage effector functions by lipoarabinomannan from mycobacterial strains with different degrees of virulence. *Infect. Immun.* **61** (1993) 4173–4181.
- BARNES, P. F., CHATTERJEE, D., ABRAMS, J. S., LU, S., WANG, E., YAMAMURA, M., BRENNAN, P. J. and MODLIN, R. L. Cytokine production induced by *Mycobacterium tuberculosis* lipoarabinomannan; relationship to chemical structure. *J. Immunol.* **149** (1992) 541–547.
- BARNES, P. F., CHATTERJEE, D., BRENNAN, P. J., REA, T. H. and MODLIN, R. L. Tumor necrosis factor production in patients with leprosy. *Infect. Immun.* **60** (1992) 1441–1446.
- BERMUDEZ, L. E. Production of transforming growth factor- β by *Mycobacterium avium*-infected human macrophages is associated with unresponsiveness to IFN- γ . *J. Immunol.* **150** (1993) 1838–1845.
- BRADBURY, M. G. and MORENO, C. Effect of lipoarabinomannan and mycobacteria on tumor necrosis factor production by different populations of murine macrophages. *Clin. Exp. Immunol.* **94** (1993) 57–63.
- CHATTERJEE, D., ROBERTS, A. D., LOWELL, K., BRENNAN, P. J. and ORME, I. M. Structural basis of capacity of lipoarabinomannan to induce secretion of tumor necrosis factor. *Infect. Immun.* **60** (1992) 1249–1253.
- DAHL, K. E., SHIRATSUCHI, H., HAMILTON, B. D., ELLNER, J. J. and TOOSI, Z. Selective induction of transforming growth factor β in human monocytes by lipoarabinomannan of *Mycobacterium tuberculosis*. *Infect. Immun.* **64** (1996) 399–405.
- DE WAAL MALEFYT, R., YSSEL, H., RONCAROLO, M.-G., SPITS, H. and VRIES, J. E. Interleukin-10. *Curr. Opin. Immunol.* **4** (1992) 314–320.
- DECKER, T., LOHMANN-MATTHES, M.-L. and GIFFORD, G. E. Cell-associated tumor necrosis factor (TNF) as a killing mechanism of activated cytotoxic macrophages. *J. Immunol.* **138** (1987) 957–962.
- DUSTIN, M. L., ROTHLEIN, R., BHAN, A. K., DINARELLO, C. A. and SPRINGER, T. A. Induction by IL 1 and interferon- γ : tissue distribution, biochemistry, and function of a natural adherence molecule (ICAM-1). *J. Immunol.* **137** (1986) 245–254.
- FRANZBLAU, S. G. Drug susceptibility testing of *Mycobacterium leprae* in the BACTEC 460 System. *Antimicrob. Agents Chemother.* **33** (1989) 2115–2117.
- HAGERTY, D. T. Intercellular adhesion molecule-1 is necessary but not sufficient to activate CD4⁺ T cells. *J. Immunol.* **156** (1996) 3652–3659.
- KRIEGLER, M., PEREZ, C., DEFAY, K., ALBERT, I. and LU, S. D. A novel form of TNF/cachectin is a surface cytotoxic transmembrane protein: ramifications for the complex physiology of TNF. *Cell* **53** (1988) 45–53.
- LOPEZ RAMIREZ, G. M., ROM, W. N., CIOTOLI, C., TALBOT, A., MARTINIUK, F., CRONSTEN, B. and REIBMAN, J. *Mycobacterium tuberculosis* alters expression of adhesion molecules on monocytic cells. *Infect. Immun.* **62** (1994) 2515–2520.
- MARLIN, S. D. and SPRINGER, T. A. Purified intercellular adhesion molecule-1 (ICAM-1) is a ligand for lymphocyte function-associated antigen-1 (LFA-1). *Cell* **51** (1987) 813–819.
- MAW, W. W., TOMIOKA, H., SATO, K. and SAITO, H. The expression of ICAM-1 on macrophages stimulated with *Mycobacterium avium* complex and its control by some regulatory cytokines. *Kekkaku* **71** (1996) 561–567.
- MONCADA, B., TORRES-ALVAREZ, M. B., GONZALEZ-AMARO, R., HUENTES-AHUMADA, C., BARANDA, L., DELGADO, S. P. and GARCIA, R. Lack of expression of intercellular adhesion molecules ICAM-1 in lepromatous leprosy patients. *Int. J. Lepr.* **61** (1993) 581–585.
- MORENO, C., TAVERNE, J., MEHLERT, A., BATE, C. A., BREALEY, R. J., MEAGER, A., ROOK, G. A. and PLAYFAIR, J. H. Lipoarabinomannan from *Mycobacterium tuberculosis* induced the production

- of tumor necrosis factor from human and murine macrophages. *Clin. Exp. Immunol.* **76** (1986) 240–245.
19. MURRAY, P. J., WANG, L., ONUFRYK, C., TEPPER, R. I. and YOUNG, R. A. T cell-derived IL-10 antagonizes macrophage function in mycobacterial infection. *J. Immunol.* **158** (1997) 315–321.
 20. ROTHLEIN, R., CZAJKOWSKI, M., O'NEILL, M. M., MARLIN, S. D., MAINOLFI, E. and MERLUZZI, V. J. Induction of intercellular adhesion molecule 1 on primary and continuous cell lines by pro-inflammatory cytokines. *J. Immunol.* **141** (1988) 1665–1669.
 21. SAITO, H., TOMIOKA, H. and SATO, K. Purification of *M. leprae* with special reference to the effects of purified *M. leprae* vaccines on host macrophage cell functions. *Jpn. J. Lepr.* **56** (1987) 101–109.
 22. SEYLER, I., APPEL, M., DEVISSAGGUET, J.-P., LEGRAND, P. and BARRATT, G. Modulation of nitric oxide production in RAW 264.7 cells by transforming growth factor-beta and interleukin-10: differential effects on free and encapsulated immunomodulator. *J. Leukoc. Biol.* **62** (1997) 374–380.
 23. SHRIKANT, P., WEBER, E., JILLING, T. and BENVENISTE, E. N. Intercellular adhesion molecule-1 gene expression by glial cells. Differential mechanisms of inhibition by IL-10 and IL-6. *J. Immunol.* **155** (1995) 1489–1501.
 24. SPRINGER, T. A. Adhesion receptors of the immune system. *Nature* **346** (1990) 425–434.
 25. SULLIVAN, L., SANO, S., PIRMEZ, C., SALGAME, P., MUELLER, C., HOFMAN, F., UYEMURA, K., REA, T. H., BLOOM, B. R. and MODLIN, R. L. Expression of adhesion molecules in leprosy lesions. *Infect. Immun.* **59** (1991) 4154–4160.
 26. TOMIOKA, H., SAITO, H. and SATO, K. Evaluation of BACTEC 460 TB System for measurement of *in vitro* anti-*Mycobacterium leprae* activity of various antimicrobials. *Jpn. J. Lepr.* **61** (1992) 157–164.
 27. TOMIOKA, H., SATO, K., MAW, W. W. and SAITO, H. The role of tumor necrosis factor, interferon- γ , transforming growth factor- β , and nitric oxide in the expression of immunosuppressive functions of splenic macrophages induced by *Mycobacterium avium* complex infection. *J. Leukoc. Biol.* **58** (1995) 704–712.
 28. TOMIOKA, H., SATO, K., SHIMIZU, T., SANO, C., AKAKI, T., SAITO, H., FUJII, K. and HIDAKA, T. Effects of benzoxazinorifamycin KRM-1648 on cytokine production at sites of *Mycobacterium avium* complex infection induced in mice. *Antimicrob. Agents Chemother.* **41** (1997) 357–362.
 29. TOOSI, Z., YOUNG, T. G., AVERILL, L. E., HAMILTON, B. D., SHIRATSUCHI, H. and ELLNER, J. J. Induction of transforming growth factor β 1 by purified protein derivative of *Mycobacterium tuberculosis*. *Infect. Immun.* **63** (1995) 224–228.
 30. VAN SEVENTER, G. A., SHIMIZU, Y., HORGAN, K. J. and SHAW, S. The LFA-1 ligand ICAM-1 provides an important costimulatory signal for T cell receptor-mediated activation of resting T cells. *J. Immunol.* **144** (1990) 4579–4586.
 31. WAHL, S. M. Transforming growth factor beta (TGF- β) in inflammation: a cause and a cure. *J. Clin. Immunol.* **12** (1992) 61–73.
 32. WILLEMS, F., MARCHANT, A., DELVILLE, J.-P., GERARD, C., DELAUX, A., VELU, T., DE BORE, M. and GOLDMAN, M. Interleukin-10 inhibits B7 and intercellular adhesion molecule-1 expression on human monocytes. *Eur. J. Immunol.* **24** (1994) 1007–1009.
 33. XIAO, B.-G., ZHANG, G.-X., MA, C.-G. and LINK, H. Transforming growth factor-beta 1 (TGF- β 1)-mediated inhibition of glial cell proliferation and downregulation of intercellular adhesion molecule-1 (ICAM-1) are interrupted by interferon-gamma (IFN- γ). *Clin. Exp. Immunol.* **103** (1996) 475–481.